

## Diclofenac residues as the cause of vulture population decline in Pakistan

J. Lindsay Oaks<sup>1</sup>, Martin Gilbert<sup>2</sup>, Munir Z. Virani<sup>2</sup>, Richard T. Watson<sup>2</sup>, Carol U. Meteyer<sup>3</sup>, Bruce A. Rideout<sup>4</sup>, H. L. Shivaprasad<sup>5</sup>, Shakeel Ahmed<sup>6</sup>, Muhammad Jamshed Iqbal Chaudhry<sup>6</sup>, Muhammad Arshad<sup>6</sup>, Shahid Mahmood<sup>6</sup>, Ahmad Ali<sup>6</sup> & Aleem Ahmed Khan<sup>6</sup>

<sup>1</sup>Department of Veterinary Microbiology and Pathology, Washington State University, Pullman, Washington 99164-7040, USA

<sup>2</sup>The Peregrine Fund, 5668 West Flying Hawk Lane, Boise, Idaho 83709, USA

<sup>3</sup>USGS-National Wildlife Health Center, 6006 Schroeder Road, Madison, Wisconsin 53711-6223, USA

<sup>4</sup>Center for Reproduction of Endangered Species, Zoological Society of San Diego, PO Box 120551, San Diego, California 92112, USA

<sup>5</sup>California Animal Health and Food Safety Laboratory System—Fresno Branch, University of California at Davis, 2789 S. Orange Avenue, Fresno, California 93725, USA

<sup>6</sup>Zoology Division, Institute of Pure and Applied Biology, Bahauddin Zakariya University, Multan, Pakistan

The Oriental white-backed vulture (OWBV; *Gyps bengalensis*) was once one of the most common raptors in the Indian subcontinent<sup>1</sup>. A population decline of >95%, starting in the 1990s, was first noted at Keoladeo National Park, India<sup>2</sup>. Since then, catastrophic declines, also involving *Gyps indicus* and *Gyps tenuirostris*, have continued to be reported across the subcontinent<sup>3</sup>. Consequently these vultures are now listed as critically endangered by BirdLife International<sup>4</sup>. In 2000, the Peregrine Fund initiated its Asian Vulture Crisis Project with the Ornithological Society of Pakistan, establishing study sites at 16 OWBV colonies in the Kasur, Khanewal and Muzaffargarh–Layyah Districts of Pakistan to measure mortality at over 2,400 active nest sites<sup>5</sup>. Between 2000 and 2003, high annual adult and subadult mortality (5–86%) and resulting population declines (34–95%) (ref. 5 and M.G., manuscript in preparation) were associated with renal failure and visceral gout. Here, we provide results that directly correlate residues of the anti-inflammatory drug diclofenac with renal failure. Diclofenac residues and renal disease were reproduced experimentally in OWBVs by direct oral exposure and through feeding vultures diclofenac-treated livestock. We propose that residues of veterinary diclofenac are responsible for the OWBV decline.

Between 2000 and 2002 we performed gross post-mortem examinations on 259 adult and subadult OWBVs, of which 219 (85%) had grossly apparent urate deposits, characteristic of visceral gout, on the surface of internal organs (Fig. 1). Visceral gout in birds is most commonly the result of renal failure leading to hyperuricaemia and the deposition of uric acid on and within the internal organs, and can be caused by degenerative, metabolic, infectious, or toxic diseases<sup>6</sup>. To verify renal disease, and to determine the cause, detailed necropsies and diagnostic testing were performed on a subset of 42 OWBVs (33 adults and 9 juveniles; 28 with visceral gout and 14 without visceral gout) that were found within approximately 24 h of death. The remaining OWBVs were significantly decomposed and thus were unsuitable for diagnostic evaluation, although the grossly apparent and characteristic lesions of visceral gout allowed the presence or absence of the disease to be determined. Of the 14 OWBVs without visceral gout, we determined the cause of death in 8 (57%), which included trauma, intestinal foreign bodies, lead poisoning, organophosphate poisoning and gun-shot (Supplementary Information). We were unable to determine the cause of death in the remaining six vultures. Among the 28 OWBVs with

visceral gout, only one (4%) had an identifiable disease (infection with *Mycobacterium avium*) in addition to visceral gout. All but two of the visceral gout cases were in good physical condition based on the subjective assessment of normal pectoral muscle mass and adequate body fat. In all of the OWBVs with visceral gout, the only significant histopathological lesion was severe, acute renal tubular necrosis and uric acid crystal formation in the kidneys and other tissues (detailed histopathological description is provided in Supplementary Information). Fibrosis or other changes indicating chronic renal disease were not present. Inflammatory lesions indicating an infectious disease were not consistently present in any of the visceral gout cases.

These findings were most compatible with acute renal failure due to a toxic cause. Additional testing eliminated known causes of avian renal disease, including toxic concentrations of cadmium, lead (39 vultures) and mercury (37 vultures)<sup>7</sup> (Supplementary Tables 1 and 2), as well as infection with the avian influenza<sup>8</sup>, infectious bronchitis<sup>9</sup> (21 vultures) and West Nile viruses<sup>10</sup> (13 vultures) (Supplementary Table 3). Testing was also negative for toxic concentrations of arsenic, and for toxic or deficient concentrations of copper, iron, manganese, molybdenum and zinc (39 vultures) (Supplementary Tables 1 and 2). Carbamate and organophosphate pesticides (34 vultures), organochlorine pesticides and polychlorinated biphenyls (13 vultures) were either not detected or were detected at below toxic concentrations (Supplementary Tables 4, 5 and 6). We failed to isolate any viruses from the kidney,

Table 1 Diclofenac residue testing results in kidney samples from wild OWBVs with and without renal failure

Case no.	Date	Site	Age	Gout	Diagnosis	Diclofenac residues ( $\mu\text{g g}^{-1}$ )
33	2001	KS	Juv	Yes	None	0.051
74	2002	KH	Ad	Yes	None	0.054
16	2001	KS	Ad	Yes	None	0.060
53	2002	KH	Ad	Yes	None	0.064
60	2002	ML	Ad	Yes	None	0.064 (0.076)
39	2001	KH	Ad	Yes	None	0.077
69	2002	ML	Ad	Yes	None	0.079
57	2002	KH	Ad	Yes	None	0.080
40	2002	ML	Ad	Yes	None	0.091
20	2001	KS	Ad	Yes	None	0.097
15	2001	KS	Ad	Yes	None	0.099
35	2001	KS	Ad	Yes	None	0.106 (0.077)
41	2002	KH	Ad	Yes	None	0.109
75	2002	KH	Ad	Yes	None	0.114
55	2002	ML	Ad	Yes	None	0.124
54	2002	ML	Ad	Yes	None	0.177
71	2002	KH	Ad	Yes	None	0.179
61	2002	ML	Ad	Yes	None	0.186
42	2002	KH	Ad	Yes	None	0.199
44	2002	KH	Ad	Yes	None	0.233
4	2000	KS	Ad	Yes	<i>M. avium</i> infection	0.450
38	2001	ML	Ad	Yes	None	0.451
59	2002	ML	Ad	Yes	None	0.504
45	2002	ML	Ad	Yes	None	0.642 (0.197)
56	2002	KH	Ad	Yes	None	0.643
2	2000	LH	Ad	No	Wire collision	BDL
3	2000	CW	Juv	No	Hit by car	BDL
12	2001	ML	Ad	No	None	BDL
14	2001	KS	Ad	No	Lead poisoning	BDL
28	2001	KS	Ad	No	Normal (trapped)	BDL (BDL)
31	2001	KH	Juv	No	Fell from nest	BDL
46	2002	KH	Juv	No	Fractured tibia	BDL
47	2002	KH	Juv	No	Intestinal foreign body	BDL
49	2002	KF	Ad	No	None	BDL
50	2002	ML	Juv	No	None	BDL
51	2002	KH	Juv	No	None	BDL
52	2002	ML	Juv	No	None	BDL
58	2002	ML	Ad	No	Organophosphate	BDL (BDL)

BDL indicates 'below detection limit' of diclofenac assay (0.005–0.01  $\mu\text{g g}^{-1}$ ). Results in parentheses are from the Toxicology Laboratory at the University of Pennsylvania New Bolton Center, which were performed as independent verification. This additional testing also did not detect acetaminophen (0.05), flunixin (0.05), ibuprofen (0.50), phenylbutazone (0.10), oxyphenbutazone (0.05), indomethacin (0.05), ketoprofen (0.25), mefenamic acid (0.50), salicylic acid (1.0), tolmetin (0.05), or naproxen (1.0) (detection limits, in parentheses, are in  $\mu\text{g g}^{-1}$ ). Ad, adult; CW, Chichawatni; Juv, juvenile; KF, Katora Forest; KH, Khanewal district study site; KS, Kasur district study site; LH, City of Lahore; ML, Muzaffargarh–Layyah district study site.

spleen, lung or intestine using at least four passages on chicken embryo fibroblast, duck embryo fibroblast, peregrine falcon (*Falco peregrinus*) embryo fibroblast and Vero cell cultures for 13 vultures (Supplementary Table 7).

The primary food source for OWBVs in Pakistan is dead domestic livestock. Therefore, we hypothesized that ingested veterinary pharmaceuticals might be responsible for renal disease in the scavenging birds. We conducted a survey of 74 veterinarians and veterinary pharmaceutical retailers in the region to identify drugs that were known to be both toxic to kidneys and absorbed orally. The only drug identified in the survey that met these criteria was diclofenac, a non-steroidal anti-inflammatory drug (NSAID) used as an analgesic, anti-inflammatory and antipyretic<sup>11</sup>. Liquid chromatography and mass spectroscopy detected diclofenac residue concentrations of 0.051–0.643  $\mu\text{g g}^{-1}$  in the kidneys of 25 out of 25 (100%) vultures that died of renal failure, and in 0 out of 13 (0%) vultures that died from other causes (Table 1). The association between renal failure and diclofenac residues was very highly significant ( $\chi^2 = 38$ , 1 degree of freedom,  $P = 0.0$ ). Independent testing on three positive and two negative samples verified these results, and did not detect the presence of any other NSAID (Table 1). Diclofenac residues were found in vultures collected between 2000 and 2002, and at study sites separated by up to 350 km.

Although there are no reports of the use of diclofenac in birds, other NSAIDs such as indomethacin<sup>12</sup>, flunixin meglumide<sup>13</sup> and ketoprofen (M. Busch & B.A.R., personal communication) may cause renal disease in chickens, cranes and quail, and in African white-backed vultures (*Gyps africanus*), respectively. NSAIDs, including diclofenac, are also recognized as causing renal disease in mammals<sup>11,14</sup>. To verify the renal toxicity of diclofenac in OWBVs, we administered single oral doses of veterinary diclofenac to four captive, non-releasable juvenile OWBVs. Two were given the mammalian label dose of 2.5  $\text{mg kg}^{-1}$  (high dose) and two were given 0.25  $\text{mg kg}^{-1}$  (low dose). Both of the high-dose and one of the low-dose vultures died as a result of renal failure and visceral gout within 36–58 h after administration, and all had the same microscopic renal lesions as the field cases. Plasma samples collected 1 h after administration for one high-dose and one low-dose vulture indicated normal uric acid levels<sup>15</sup> (53 and 33  $\text{mg l}^{-1}$ , respectively). By 24 h after administration, both vultures had developed hyper-

uricaemia (775  $\text{mg l}^{-1}$  for both). Diclofenac residue concentrations of 0.29, 1.1 and 0.16  $\mu\text{g g}^{-1}$  were present in the kidneys of the two high-dose vultures and the one low-dose vulture that died as a result of renal failure. Although the other low-dose vulture developed hyperuricaemia (138  $\text{mg l}^{-1}$  at 1 h and 654  $\text{mg l}^{-1}$  at 24 h after administration), it remained clinically normal 4 weeks after administration, and did not have microscopic renal lesions or detectable diclofenac residues at necropsy. Two untreated control birds that were fed the same diet also did not have renal lesions or detectable diclofenac residues at necropsy.

The most probable source of diclofenac exposure is the consumption of treated livestock. In Pakistan, veterinary diclofenac is widely marketed over the counter by multiple companies for treating all types of hooved livestock. A second survey of 84 drug retailers and veterinarians in nine districts in the Punjab province (Dera Ghazi Khan, Kasur, Khanewal, Layyah, Multan, Muzaffargarh, Pakpattan, Rajanpur, Sahiwal) and one district in the Sindh province (Hyderabad) found that all 84 sold diclofenac; 77 sold it daily, and 71 reported that it had become available within the last 5 years. Furthermore, livestock that die of disease or injury are typically left for scavengers to remove.

To verify that carcasses of treated livestock contain sufficient diclofenac concentrations to cause renal failure and death in the vultures that scavenge upon them, ten juvenile OWBVs were fed meat from a buffalo or goat that was injected intramuscularly with 2.5  $\text{mg kg}^{-1}$  veterinary diclofenac once daily for 3 days (per label instructions) and that were slaughtered 4 h hours after the last injection. Resulting diclofenac residue concentrations in the buffalo kidney, liver and muscle were 5.7, 1.5 and 0.76  $\mu\text{g g}^{-1}$ , respectively, and in the goat kidney, liver and muscle they were 0.94, 0.22 and 0.19  $\mu\text{g g}^{-1}$ , respectively. Ten more OWBVs were fed buffalo meat that contained 6.4  $\mu\text{g g}^{-1}$  of diclofenac. On the basis of the product of diclofenac concentration in the meat and the amount of food consumed, eight OWBVs received doses of 0.005–0.3  $\text{mg kg}^{-1}$ . Two of these vultures died from renal failure at 4 and 6 days after exposure, and at necropsy they had diclofenac residue concentrations of 0.07 and 0.38  $\mu\text{g g}^{-1}$  in the kidney. A necropsy was carried out on one surviving, clinically normal vulture at 8 days after exposure, at which time it did not have renal lesions or detectable diclofenac residues. The other five surviving vultures remain clinically normal at approximately 6 months after exposure. Two



**Figure 1** Necropsy photograph of the abdominal cavity of an OWBV with visceral gout, showing uric acid precipitates on the serosal surfaces of the liver.

OWBVs received doses of 0.5–0.6 mg kg<sup>-1</sup>. One of these died from renal failure 1 day after exposure, and at necropsy it had a diclofenac residue concentration of 0.25 µg g<sup>-1</sup> in the kidney. The other surviving, clinically normal vulture had a necropsy performed on it at 8 days after exposure, at which time it did not have renal lesions or detectable diclofenac residues. Ten OWBVs received doses of 0.8–1.0 mg kg<sup>-1</sup>. All ten of these vultures died from renal failure. Kidney diclofenac residue concentrations of 0.120–0.906 µg g<sup>-1</sup> were present in six out of six of these vultures tested. These ten vultures were negative for other recognized avian nephrotoxins (Supplementary Information). All 13 vultures that died from renal failure after consuming meat from diclofenac-treated animals had the same histopathological renal lesions as the vultures that were exposed directly with diclofenac, and as the field cases with visceral gout. Six control vultures that were fed untreated goat (meat that did not contain detectable diclofenac residues) remained clinically normal 8 and 25 days into the experiment. Two were necropsied at day eight, and they did not have renal lesions or detectable diclofenac residues. These results are summarized in Table 2. In addition, all of the experimental OWBVs were held in captivity, and were clinically normal for 30–75 days before diclofenac exposure, indicating that visceral gout was not incidentally related to captivity. The combined mortality rate of 13 out of 20 (65%) in the exposed vultures and 0 out of 6 (0%) in the control vultures indicates a statistically highly significant relationship between renal failure and exposure to diclofenac ( $\chi^2 = 7.8$ , 1 degree of freedom,  $P = 0.005$ ). When the direct and food-borne exposure levels were combined, a statistically highly significant ( $\chi^2 = 12.3$ , 3 degrees of freedom,  $P = 0.0064$ ) dose-dependence for diclofenac toxicity can also be demonstrated.

Alternatively, wild OWBVs might be exposed to diclofenac through contaminated water sources<sup>16</sup>; however, because NSAID-associated renal failure in birds appears to be acute and dose-dependent<sup>12,13</sup>, the very low water concentrations (on the scale of ng l<sup>-1</sup>)<sup>16</sup> would be unlikely to cause toxicity. Also, the rapid and complete elimination of most NSAIDs in mammals<sup>14</sup> and birds<sup>17</sup> makes bioaccumulation in either the treated animals or in the scavengers that feed on them unlikely. The lack of detectable diclofenac residues in the kidneys of any unaffected wild OWBV

and in the kidneys of the surviving experimental vultures also suggest that bioaccumulation does not occur.

The potential ecologically toxic effects of human and veterinary pharmaceuticals are a growing concern<sup>16,18</sup>. Sporadic poisonings of scavenging birds by organophosphate pesticides and barbiturates used in livestock have been documented previously<sup>19,20</sup>. Anthelmintics may be shed in livestock faeces and may kill ecologically important invertebrates<sup>21</sup>. However, in stark contrast to the current situation with the OWBV, pharmaceutical residues have not been implicated previously as a cause of major ecological damage. The identification of diclofenac as the cause of the OWBV decline in Pakistan provides an opportunity for conservation intervention. The high rate of visceral-gout-associated vulture mortality in India<sup>3,22,23</sup> as well as the widespread use of veterinary diclofenac in India (R. Risebrough, personal communication) suggests strongly that diclofenac may also be responsible for vulture declines in the rest of the Indian subcontinent wherever diclofenac is used for the treatment of livestock. □

Methods

Histopathology

Tissue samples were fixed in 10% neutral-buffered formalin, and 5-µm sections were cut and stained routinely with haematoxylin and eosin.

Toxicology

Testing for arsenic, cadmium, copper, iron, lead, manganese, mercury, molybdenum and zinc was performed by the inductively coupled plasma method. Organophosphate and carbamate testing included measuring brain acetylcholinesterase levels<sup>24</sup> as an indirect indicator of exposure<sup>25</sup>, and direct detection by liquid chromatography and mass spectroscopy on liver, brain or stomach contents. Organochlorine pesticide and polychlorinated biphenyl testing was performed by liquid chromatography and mass spectroscopy. Reported concentrations are based on the wet weight of the samples.

Diclofenac analysis

Kidney samples (0.5 g) were homogenized in 4 ml of acetonitrile and centrifuged to pellet debris. Three millilitres of the supernatants were then passed through solid-phase extraction cartridges (Waters Sep-Pak Plus T-C18) at 1 ml min<sup>-1</sup>. The cartridges were then washed with an additional 3 ml of acetonitrile, the eluates pooled, and concentrated to final volumes of 1 ml. For samples that weighed less than 0.5 g, the protocol was adjusted proportionally for the smaller weights.

Diclofenac was detected and quantified by high-performance liquid chromatography and mass spectroscopy (Agilent 1100 series equipped with a Water's Xterra MS C18 (3.9 mm × 150 mm, 5 µm) and guard column (3.9 mm × 20 mm, 5 µm)). Diclofenac standard (Sigma, D6899) was dissolved into 1:1 (v/v) acetonitrile:water at 1,040 µg ml<sup>-1</sup>, diluted to a final concentration of 10.4 µg ml<sup>-1</sup>, and this stock was used to prepare calibration standards of 0.104, 0.052, 0.026, 0.0104 and 0.0052 µg ml<sup>-1</sup> in acetonitrile, which generated a linear calibration curve with R<sup>2</sup> values equal to 0.999. Samples and standards (20 µl) were subjected to a binary gradient elution profile, which consisted of 0.1% acetic acid in water (solution A) and 100% acetonitrile (solution B) as follows: starting conditions 75% A/25% B for 0.1 min, a 15-min linear gradient from 75% A/25% B to 5% A/95% B, followed by a 5-min column-wash step in 5% A/95% B, and a 10-min re-equilibration step with 75% A/25% B before the next injection. Flow rate was 0.7 ml min<sup>-1</sup> and the column temperature was 40 °C. The mass spectrometer (Agilent 1946D) was equipped with an electrospray ionization inlet and mass spectra were acquired in the negative ion mode. The mass spectrometer was set to selectively monitor for mass ions 294 and 296 m/z of which mass 294 m/z, the deprotonated molecular mass, was used to quantify the diclofenac residues. The mean recovery from spiked avian kidney tissue was approximately 70%, and this value was used to calculate the final sample concentration of diclofenac.

Virus isolation

Virus isolation in cell culture was performed in primary avian cell cultures prepared by the method of ref. 26 from chicken, duck and peregrine falcon embryos, and in Vero cells. Tissue samples were prepared by homogenization in Eagle's minimal essential medium with penicillin, streptomycin and amphotericin B, centrifuged to remove debris, filtered through 0.45-µm filters, and added to the cells. Cell cultures were observed daily for cytopathology. At the end of each passage (5–8 days), the cells and media were aspirated, stored at -80 °C, rapidly thawed, and added to new cells for the next passage.

Polymerase chain reaction assays for infectious agents

RNA and DNA were extracted from 0.1–0.3 g of tissue with a commercial RNA extraction kit (Trizol Reagent; BRL Life Technologies) or a commercial DNA extraction kit (Puregene; Gentra Systems) as per the manufacturer's instructions. In addition to positive and negative control RNA or DNA for each agent, intact sample RNA and DNA were verified by demonstrating the presence of avian cellular β-actin gene RNA or DNA, and the absence of nonspecific inhibitors was verified by spiking samples with positive control RNA or DNA.

Table 2 Results of experimentally feeding meat from diclofenac-treated animals to OWBVs

Vulture no	Exposure (mg kg <sup>-1</sup> )	Result	Diclofenac residues (µg g <sup>-1</sup> )
11	0.007	Gout	0.38
12	0.025	Survived	ND
8	0.027	Survived	ND
10	0.028	Survived	ND
7	0.029	Survived	ND
9	0.029	Survived	ND
D	0.140	Gout	0.07
F	0.240	Survived	BDL (8 days)
H	0.550	Gout	0.25
B	0.600	Survived	BDL (8 days)
64	0.820	Gout	0.54
65	0.820	Gout	ND
62	0.840	Gout	0.22
63	0.840	Gout	0.12
70	0.860	Gout	0.74
67	0.860	Gout	ND
68	0.860	Gout	ND
66	0.880	Gout	0.26
73	0.910	Gout	0.91
72	0.940	Gout	ND
1	None (control)	Survived	ND
2	None (control)	Survived	ND
3	None (control)	Survived	ND
4	None (control)	Survived	ND
cc1	None (control)	Survived	BDL (8 days)
cc2	None (control)	Survived	BDL (8 days)

Values in parentheses in the last column indicate time after exposure. OWBVs that died and displayed gross characteristics of visceral gout on necropsy are labelled as 'Gout' in the third column. BDL, below detection limit of diclofenac assay (0.005–0.01 µg g<sup>-1</sup>); ND, analysis not done.

The polymerase chain reaction with the reverse transcription (RT-PCR) procedure for infectious bronchitis virus was as published in ref. 27—the positive control for this assay was the Arkansas strain. The RT-PCR procedure for avian influenza was as published in ref. 28—the positive control for this assay was viral RNA obtained from the USDA National Veterinary Services Laboratory, Ames, Iowa. The RT-PCR procedure for West Nile virus was as published in ref. 29—the positive control for this assay was viral RNA obtained from the USDA National Veterinary Services Laboratory, Ames, Iowa.

## Identification of *M. avium*

Acid-fast bacilli were detected with Ziehl-Neelsen stains of tissue impression smears. *Mycobacterium avium* was identified by PCR amplification and restriction enzyme polymorphisms in the heat-shock protein gene<sup>30</sup>.

## Uric acid measurement

Uric acid levels were measured in heparinized plasma with a colorimetric assay for the production of hydrogen peroxide by uricase.

## Animal subjects

Animals were maintained and experiments performed in accordance with the Animal Health and Welfare Regulations of Bahauddin Zakariya University, Pakistan.

Received 14 September; accepted 23 December 2003; doi:10.1038/nature02317.

Published online 28 January 2004.

1. Ali, S. & Ripley, S. D. *Handbook of the Birds of India and Pakistan Together With Those of Nepal, Bhutan, and Ceylon I. Divers to Hawks* (Oxford Univ. Press, Oxford, 1968).
2. Prakash, V. Status of vultures in Keoladeo National Park, Bharatpur, Rajasthan, with special reference to population crash in *Gyps* species. *J. Bombay Nat. Hist. Soc.* **96**, 365–378 (1999).
3. Pain, D. *et al.* Causes and effects of temporospatial declines of *Gyps* vultures in Asia. *Conserv. Biol.* **17**, 661–671 (2003).
4. BirdLife International. *Threatened Birds of Asia: the BirdLife International Red Data Book* (BirdLife International, Cambridge, 2001).
5. Gilbert, M. *et al.* Breeding and mortality of Oriental White-backed vulture *Gyps bengalensis* in Punjab Province, Pakistan. *Bird Conserv. Int.* **12**, 311–326 (2002).
6. Crespo, R. & Shivaprasad, H. L. in *Diseases of Poultry* 11th edn (ed. Saif, Y. M.) 1085–1087 (Iowa State Press, Ames, 2003).
7. Beyer, W., Heinz, G. & Redmon-Norwood, A. (eds) *Environmental Contaminants in Wildlife. Interpreting Tissue Concentrations* Ch. 10, 11, 14, 17 (CRC Press, Boca Raton, 1996).
8. Swayne, D. E. & Slemons, R. D. Comparative pathology of a chicken-origin and two-duck origin influenza virus isolates in chickens: the effect of route of inoculation. *Vet. Pathol.* **31**, 237–245 (1994).
9. Ziegler, A. F. *et al.* Nephropathogenic infectious bronchitis in Pennsylvania chickens 1997–2000. *Avian Dis.* **46**, 847–858 (2002).
10. Steele, K. E. *et al.* Pathology of fatal West Nile virus infections in native and exotic birds during the 1999 outbreak in New York City, New York. *Vet. Pathol.* **37**, 208–224 (2000).
11. Todd, P. A. & Sorokin, E. M. Diclofenac sodium. A reappraisal of its pharmacodynamic and pharmacokinetic properties, and therapeutic efficacy. *Drugs* **35**, 244–285 (1988).
12. Nys, Y. & Rzasa, J. Increase in uricemia induced by indomethacin in hens or chickens. *C. R. Séances Acad. Sci. III* **296**, 401–404 (1983).
13. Paul-Murphy, J. & Ludders, J. W. Avian analgesia. *Vet. Clin. N. Am. Exotic Anim. Pract.* **4**, 35–45 (2001).
14. Murray, M. D. & Brater, D. C. Renal toxicity of the nonsteroidal anti-inflammatory drugs. *Annu. Rev. Pharmacol. Toxicol.* **33**, 435–465 (1993).
15. Gatome, C. W. *Haematology and Blood Biochemistry in Free-Living African White-Backed Vultures Gyps africanus in Kenya* (MSc, Univ. London, 2002).
16. Tixier, C., Singer, H. P., Oellers, S. & Muller, S. R. Occurrence and fate of carbamazepine, clofibrate, diclofenac, ibuprofen, ketoprofen, and naproxen in surface waters. *Environ. Sci. Technol.* **37**, 1061–1068 (2003).

17. Baert, K. & De Backer, P. Comparative pharmacokinetics of three non-steroidal anti-inflammatory drugs in five bird species. *Comp. Biochem. Physiol. C* **134**, 25–33 (2003).
18. Daughton, C. G. & Ternes, T. A. Pharmaceuticals and personal care products in the environment: agents of subtle change. *Environ. Health Perspect.* **107**, 907–938 (1999).
19. Henny, C. J., Blus, L. J., Kolbe, E. J. & Fitzner, R. E. Organophosphate insecticide (famphur) topically applied to cattle kills magpies and hawks. *J. Wildl. Mgmt* **49**, 648–658 (1985).
20. Langelier, K. M. in *Raptor Biomedicine* (eds Redig, P. T., Cooper, J. E., Rempke, J. D. & Hunter, D. B.) 231–232 (Univ. Minnesota, Minneapolis, 1993).
21. McKellar, Q. A. Ecotoxicology and residues of anthelmintic compounds. *Vet. Parasitol.* **72**, 413–435 (1997).
22. Cunningham, A. A. *et al.* Indian vultures: victims of an infectious disease epidemic? *Anim. Conserv.* **6**, 189–197 (2003).
23. Mishra, S. K. *et al.* Vulture mortality: pathological and microbiological investigations. *Indian J. Anim. Sci.* **72**, 283–286 (2002).
24. Ellman, G. L., Courtney, K. D., Andres, V. & Featherstone, R. M. A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochem. Pharmacol.* **7**, 88–95 (1961).
25. Hill, E. F. & Fleming, W. J. Anticholinesterase poisoning of birds: field monitoring and diagnosis of acute poisoning. *Environ. Toxicol. Chem.* **1**, 27–38 (1982).
26. Dochertry, D. E. & Slota, P. G. Use of muscovy duck embryo fibroblasts for the isolation of viruses from wild birds. *J. Tissue Cult. Methods* **11**, 165–170 (1988).
27. Kwon, H. M., Jackwood, M. W. & Gelb, J. Differentiation of infectious bronchitis virus serotypes using polymerase chain reaction and restriction fragment length polymorphism analysis. *Avian Dis.* **37**, 194–202 (1993).
28. Lee, M. S., Chang, P., Shien, J., Cheng, M. & Shieh, H. K. Identification and subtyping of avian influenza viruses by reverse transcription-PCR. *J. Virol. Methods* **97**, 13–22 (2001).
29. Johnson, D. J., Ostlund, E. N., Pedersen, D. D. & Schmitt, B. J. Detection of North American West Nile virus in animal tissue by a reverse transcription-nested polymerase chain reaction assay. *Emerg. Inf. Dis.* **7**, 739–741 (2001).
30. Mendenhall, M. K. *et al.* Detection and differentiation of *Mycobacterium avium* and *Mycobacterium genavense* by polymerase chain reaction and restriction enzyme digestion analysis. *J. Vet. Diagn. Invest.* **12**, 57–60 (2000).

**Supplementary Information** accompanies the paper on [www.nature.com/nature](http://www.nature.com/nature).

**Acknowledgements** The authors thank S. Pritchard, S. Donahoe and M. Asim for technical assistance, and T. Taruscio and HEJ Laboratories (Karachi) for performing diclofenac assays. We thank V. Beasley, K. Beckman, P. Benson, T. Besser, T. Cade, K. Mealey, R. Poppenga, P. Talcott, The Ornithological Society of Pakistan, Punjab Department of Wildlife and Parks, the National Council for the Conservation of Wildlife (Islamabad), World Wildlife Fund (Pakistan), and Brigadier M. Ahmed for their cooperation. This research was conducted as part of The Peregrine Fund's Asian Vulture Crisis Project and was supported by the Gordon and Betty Moore Foundation, The Peregrine Fund, Disney Wildlife Conservation Fund, San Diego Zoological Society, and the United Nations, Ivorybill and Summit Foundations.

**Authors' contributions** J.L.O. coordinated the overall diagnostic work and performed microbiology; M.G., M.Z.V. and A.A.K. coordinated field biology and collected samples; R.T.W. supervised field biology; C.U.M., B.A.R. and H.L.S. performed pathology; and S.A., M.J.I.C., M.A. and A.A. collected biology data and samples.

**Competing interests statement** The authors declare that they have no competing financial interests.

**Correspondence** and requests for materials should be addressed to J.L.O. (loaks@vetmed.wsu.edu).